

(19)

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 088 550 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
04.04.2001 Bulletin 2001/14

(51) Int Cl.7: **A61K 31/18**, A61P 25/16,  
A61P 25/28

(21) Application number: **00308442.3**

(22) Date of filing: **27.09.2000**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(72) Inventors:  
• **Sahagan, Barbara Gail**  
Groton, Connecticut 06340 (US)  
• **Villalobos, Anabella**  
Groton, Connecticut 06340 (US)


(30) Priority: **01.10.1999 US 157083 P**

(74) Representative: **Ruddock, Kelth Stephen et al**  
**Pfizer Limited,**  
European Patent Department,  
Ramsgate Road  
Sandwich, Kent CT13 9NJ (GB)

(71) Applicant: **Pfizer Products Inc.**  
Groton, Connecticut 06340 (US)

(54) **alpha-sulfonylamino hydroxamic acid inhibitors of matrix metallo-proteinases for the treatment of peripheral or central nervous system disorders**

(57) The present invention relates to a method of using a compound of the formula (I):



wherein A, X<sup>1</sup>, X<sup>2</sup>, and Q are as defined herein, a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in the treatment of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to Alzheimer's disease, stroke/cerebral ischemia, head trauma, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, migraine, cerebral amyloid angiopathy, AIDS, age-related cognitive decline, mild cognitive impairment and prion diseases.

**EP 1 088 550 A1**

## Description

## Background of the Invention

- 5 [0001] The present invention relates to new methods of using certain  $\alpha$ -sulfonylamino hydroxamic acid inhibitors of matrix metalloproteinases in the treatment of diseases, conditions and disorders of the peripheral or central nervous system, including but not limited to Alzheimer's disease, stroke/cerebral ischemia, head trauma, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, migraine, cerebral amyloid angiopathy, AIDS, age-related cognitive decline; mild cognitive impairment and prion diseases, and pharmaceutical compositions useful therefor.
- 10 [0002] The compounds of the present invention are inhibitors of zinc metalloendopeptidases, especially those belonging to the matrix metalloproteinase (also called MMP or matrixin) and reprolysin (also known as adamalysin) sub-families of the metzincins (Rawlings, *et al.*, Methods in Enzymology, 248, 183-228 (1995) and Stocker, *et al.*, Protein Science, 4, 823-840 (1995)).
- 15 [0003] The MMP subfamily of enzymes, currently contains seventeen members (MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-10, MMP-11, MMP-12, MMP-13, MMP-14, MMP-15, MMP-16, MMP-17, MMP-18, MMP-19, MMP-20). The MMP's are most well known for their role in regulating the turn-over of extracellular matrix (ECM) proteins and as such play important roles in normal physiological processes such as reproduction, development and differentiation.
- 20 [0004] In the central nervous system, the ECM not only serves structural and adhesive functions but also stimulates intracellular signaling pathways in response to association of the matrix with cell surface proteins. (Yong *et al.*, Trends in Neuroscience, 21, 75-80 (1998)) Excessive expression of MMP's is believed to contribute to the pathogenesis of tissue destructive diseases such as arthritis, multiple sclerosis (MS) and cancer, conditions where inflammation and invasive processes play important roles. In Alzheimer's Disease (AD) and age-matched control samples, the expression of MMP's, particularly MMP-9 and MMP-2, is increased. The link between MMP's, AD and the ECM is supported by
- 25 *in vitro* and *in vivo* evidence. In clinical samples taken from stroke, MS, amyotrophic lateral sclerosis (ALS) patients, increased expression of MMP's has also been documented. In AD, astrocytes produce inflammatory mediators and ECM proteins which surround neuritic plaques.
- [0005] Like other members of the matrix metalloproteinase family, MMP-2 (72kDa type IV collagenase or Gelatinase A) and MMP-9 (92kDa type IV collagenase or Gelatinase B) are calcium-requiring, zinc containing endopeptidases which are secreted from cells in a latent pro-enzyme form (Yong *et al.*, *supra*). These MMP's attack type IV collagen, laminin and fibronectin, the major components of the ECM surrounding cerebral blood vessels. Because of the dire consequences of inappropriate or unbalanced activity, they are tightly regulated by three independent mechanisms: proenzyme activation, gene transcription and inhibition by their endogenous inhibitor TIMP-1 (Borden and Heller, Critical Reviews in Eukaryotic Gene Expression, 7, 159-178, (1997)). The expression of MMP-9 is induced by growth factors and inflammatory cytokines in an NF-KB and AP-1 dependent manner (Bond *et al.*, FEBS Letters, 435, 29-34, (1998)). MMP-2 is generally constitutively expressed; however, its mRNA can be modulated by some of the same factors which modulate MMP-9 expression (Gottschall and Deb, Neuroimmunomodulation, 3, 69-75, (1996)).
- 30 [0006] Additionally, in AD hippocampus, MMP-9 protein is increased as much as fourfold (Backstrom *et al.*, J. Neurochemistry, 58, 983-92 (1992)). The enzyme is predominantly found in its latent or proenzyme form in close proximity to extracellular amyloid plaques (Backstrom *et al.*, J. Neuroscience, 16, 7910-19 (1996)). Similar observations were made in aged beagles. In amyloid-positive samples, statistically significant increases in latent MMP-9 were seen as compared with amyloid-negative samples (Lim *et al.*, J. Neurochemistry, 68, 1606-11 (1997)).
- 35 [0007] The link between MMPs, AD and the ECM is supported by additional evidence (Perlmutter *et al.*, J. Neuroscience Res., 30, 673-81 (1991); Brandan and Inestrosa, Gen. Pharmacology, 24, 1063-8 (1993); Eikelenboom *et al.*, Virchows Archiv, 424, 421-7 (1994); Luckenbill-Edds, Brain Res. Revs., 23, 1-27 (1997)). Laminin is induced by brain injury and co-localizes with amyloid deposits in AD. In AD tissue, native human laminin was localized in large punctate, extracellular deposits which co-localize with plaques. Antibodies to the neurite-outgrowth promoting domains of laminin B2 or A chains localize to glia or capillary basement membranes, respectively. In control brains, laminin immunoreactivity is only found in capillaries (Murtomaki *et al.*, J. Neuroscience Res., 32, 261-73 (1993)). In a murine model of neurodegeneration (Chen and Strickland, Cell, 91, 917-25 (1997)), kainic acid challenged neurons secrete tPA. This initiates a cascade of proteolytic events beginning with conversion of plasminogen to plasmin and ending with degradation of laminin and subsequent death of neurons. Plasmin is a known activator of MMP-9 which could be part of this proteolytic cascade resulting in the eventual destruction of neurons.
- 40 [0008] In PC12 cells, laminin or specific laminin peptides are capable to stimulating MMP secretion and this mechanism is linked to laminin-mediated neurite outgrowth (Weeks *et al.*, Exp. Cell Res., 243, 375-82 (1998)). There has been a suggestion that these specific laminin sites may only be exposed in the basement membrane as observed in AD (Kibby *et al.*, Proc. Nat. Acad. Sci., 90, 10150-3 (1993)). Further deposition of A $\beta$  could be nucleated by these
- 55

laminin fragments which are found in neuritic plaques. Therefore, interfering with degradation of laminin could have the outcome of preserving the ECM, enhancing neuronal survival, and eliminating at least one protein which may act as a seed for nucleation of A $\beta$ .

[0009] Elevated expression of MMP-9 and MMP-2 has also been documented in stroke, MS and ALS. After focal ischemia in humans, MMP-9 is markedly elevated in the infarcted tissue at two days post-infarction and remained elevated for months. Increases in MMP-2 were subtle at 2-5 days and like MMP-9, remained marked and significant for months (Clark *et al.*, Neuroscience Letters, 238, 53-6 (1997)). Analysis of brain and spinal cord samples from ALS patients identified major bands of enzyme activity as MMP-2 and MMP-9; MMP-2 in astrocytes and MMP-9 in pyramidal neurons of the motor cortex and motor neurons of the spinal cord. Increases in MMP-9 were observed in ALS frontal and occipital cortices and spinal cord versus control samples. The high level of MMP-9 and its possible release at the synapse may destroy the structural integrity of the surrounding matrix thereby contributing to the pathogenesis of ALS. (Lim *et al.*, J. Neurochemistry, 67, 251-9 (1996)). MMP-9 is elevated in CSF of MS patients and is detected by immunohistochemistry in active and chronic lesions. In autopsied samples from normal brain, MMP-like immunoreactivity (MMP-1, -2, -3 and -9) is localized to microglia and astrocytes. In MS patient samples, MMP expression is up-regulated in these glial cells and also in perivascular macrophages that are present in active brain lesion. (Chandler *et al.*, J. Neuroimmunology, 72, 155-61 (1997); Liedtke *et al.*, Annals of Neurology, 44, 35-46 (1998).)

[0010] In addition to the foregoing, MMP's have been associated with neuronal degeneration in a number of animal models. These models can be used in an MMP inhibitor program to track inhibitor activity and predict pre-clinical efficacy. After focal ischemia in rats, MMP-9 was shown to increase in the infarcted area during the first day (Rosenberg *et al.*, J. Cerebral Blood Flow & Metabolism, 16, 360-6, (1996)). MMP-2 remained the same until 5 days after injury when it increased significantly. This time course of induction is very similar to that seen by Clark *et al.*, *supra*, in human clinical stroke samples. (Rosenberg and Navratil, Neurology, 48, 921-6 (1997)) have also shown that metalloproteinase inhibition blocks edema in intracerebral hemorrhage in the rat. A model of direct injection of MMP's into the rat brain also demonstrated neuronal loss after MMP-9 or -2, but not MMP-8, injection as well as the loss of GFAP and myelin immunoreactivity (Anthony *et al.*, J. Neuroimmunology, 87, 62-72 (1998)).

[0011] MMP-9 was detected in the CSF of mice with Experimental Autoimmune Encephelomyelitis (EAE), an animal model for MS. A hydroxamate inhibitor of MMP, GM6001, was found to suppress the development of, or reverse established, EAE (Gijbels, J. Clinical Invest., 94, 2177-82 (1994)). Based on this model, MMP inhibitors might act by preventing the influx of inflammatory cells across the basement membrane or ECM barrier that surrounds cerebral endothelium. Another hydroxamic acid-based compound which is a combined inhibitor of MMP and TNF- $\alpha$  processing/release, BB-1101 (Redford *et al.*, Brain, 120, 1895-905 (1997)), attenuates Experimental Autoimmune Neuritis (EAN), a model of Guillain-Barre syndrome.

[0012] Lastly, *in vitro* experiments suggest a role for MMP's in the development or progression of neuritic plaques. Deb and Gottschall, J. Neurochemistry, 66, 1641-7 (1996)), demonstrated that A $\beta$  induces MMP-9 and -2 expression in astrocyte and mixed hippocampal cultures. Further, MMP-9 induction by A $\beta$  in rat microglia can be inhibited by the anti-inflammatory agents dexamethasone and indomethacin (Gottschall, Neuroreport, 7, 3077-80 (1996)). This is of particular importance when considered in conjunction with clinical data which suggests that administration of anti-inflammatory drugs may slow the progression of AD (Rogers *et al.*, Arzneimittelforschung, 45, 439-42 (1995)).

[0013] Applicants now disclose a method of treatment of diseases, conditions or disorders of the peripheral and central nervous system, including but not limited to Alzheimer's disease, stroke/cerebral ischemia, head trauma, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, migraine, cerebral amyloid angiopathy, AIDS, age-related cognitive decline, mild cognitive impairment and prion diseases, comprising the administration of small molecule inhibitors of MMP-9, MMP-2 or mixed MMP inhibitors which may reduce neuronal damage and limit neuroinflammation.


[0014] However, the diseases in which inhibition of MMP's will provide therapeutic benefit include but are not limited to: arthritis (including osteoarthritis and rheumatoid arthritis), inflammatory bowel disease, Crohn's disease, emphysema, acute respiratory distress syndrome, asthma chronic obstructive pulmonary disease, Alzheimer's disease, organ transplant toxicity, cachexia, allergic reactions, allergic contact hypersensitivity, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, osteoporosis, loosening of artificial joint implants, atherosclerosis (including atherosclerotic plaque rupture), aortic aneurysm (including abdominal aortic aneurysm and brain aortic aneurysm), congestive heart failure, myocardial infarction, stroke, cerebral ischemia, head trauma, spinal cord injury, neurodegenerative disorders (acute and chronic), inflammatory and autoimmune disorders, Huntington's disease, Parkinson's disease, migraine, depression, peripheral neuropathy, pain, cerebral amyloid angiopathy, nootropic or cognition enhancement, amyotrophic lateral sclerosis, multiple sclerosis, ocular angiogenesis, corneal injury, macular degeneration, abnormal wound healing, burns, diabetes, tumor invasion, tumor growth, tumor metastasis, corneal scarring, scleritis, AIDS, sepsis, septic shock and other diseases characterized by metalloproteinase or ADAM expression. This invention also relates to methods of using the hydroxamic acid compounds described herein in the treatment of the above-identified diseases, conditions and disorders in mammals, especially humans, and to the pharmaceutical compositions

containing these compounds useful in such methods.

[0015] It is recognized that different combinations of MMP's are expressed in different pathological situations. As such inhibitors with specific selectivities for individual MMP's may be preferred for individual diseases.

## 5 Summary of the Invention


[0016] The present invention relates to a method of treating a disease or disorder of the peripheral or central nervous system, including but not limited to Alzheimer's disease, stroke/cerebral ischemia, head trauma, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, migraine, cerebral amyloid angiopathy, AIDS, age-related cognitive decline; mild cognitive impairment and prion diseases in a mammal, which comprises administering to said mammal a therapeutically effective amount of a compound of formula (I):




or the pharmaceutically acceptable salts thereof, wherein


A is H or  $-(CH_2)_n-(C=O)-Z$ ; where n is 0 to 6; and Z is hydroxy,  $(C_1-C_6)$ alkoxy or  $NR^1R^2$  wherein  $R^1$  and  $R^2$  are each independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl, piperidyl,  $(C_1-C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ arylpiperidyl,  $(C_2-C_9)$ heteroarylpiperidyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylpiperidyl,  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkylpiperidyl,  $(C_1-C_6)$ acylpiperidyl,  $(C_6-C_{10})$ aryl,  $(C_2-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryl $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl $(C_1-C_6)$ alkyl,  $R^5$   $(C_2-C_6)$ alkyl,  $(C_1-C_5)$ alkyl $(CHR^3)(C_1-C_6)$ alkyl wherein  $R^3$  is hydroxy,  $(C_1-C_6)$ acyloxy,  $(C_1-C_6)$ alkoxy, piperazino,  $(C_1-C_6)$ acylamino,  $(C_1-C_6)$ alkylthio,  $(C_6-C_{10})$ arylthio,  $(C_1-C_6)$ alkylsulfinyl,  $(C_6-C_{10})$ arylsulfinyl,  $(C_1-C_6)$ alkylsulfoxyl,  $(C_6-C_{10})$ arylsulfoxyl, amino,  $(C_1-C_6)$ alkylamino,  $((C_1-C_6)alkyl)_2$  amino,  $(C_1-C_6)$ acylpiperazino,  $(C_1-C_6)$ alkylpiperazino,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylpiperazino,  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkylpiperazino, morpholino, thiomorpholino, piperidino or pyrrolidino;  $R^4(C_1-C_6)$ alkyl,  $(C_1-C_5)$ alkyl $(CHR^4)(C_1-C_6)$ alkyl wherein  $R^4$  is piperidyl,  $(C_1-C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ arylpiperidyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylpiperidyl,  $(C_2-C_9)$ heteroarylpiperidyl or  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkylpiperidyl; and  $CH(R^5)COR^6$  wherein  $R^5$  is hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylthio $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylthio $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfinyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfinyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfonyl $(C_1-C_6)$ alkyl, hydroxy $(C_1-C_6)$ alkyl, amino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $((C_1-C_6)alkylamino)_2(C_1-C_6)$ alkyl,  $R^7R^8NCO(C_1-C_6)alkyl$  or  $R^7OCO(C_1-C_6)alkyl$  wherein  $R^7$  and  $R^8$  are each independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl and  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkyl; and  $R^6$  is  $R^9O$  or  $R^9R^{10}N$  wherein  $R^9$  and  $R^{10}$  are each independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl and  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkyl;

or  $R^1$  and  $R^2$ , or  $R^7$  and  $R^8$ , or  $R^9$  and  $R^{10}$  may be taken together to form an azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, indolinyl, isoindolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl,  $(C_1-C_6)$ acylpiperminyl,  $(C_1-C_6)$ alkylpiperazinyl,  $(C_6-C_{10})$ arylpiperazinyl,  $(C_2-C_9)$ heteroarylpiperazinyl or a bridged diazabicycloalkyl ring selected from the group consisting of:



(J<sub>4</sub>)

and

(J<sub>5</sub>)

wherein p is 1, 2 or 3;


q is 1 or 2;

r is 0 or 1;

L is hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)acyl;

X<sup>1</sup> and X<sup>2</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoromethyl, trifluoromethyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl (difluoromethylene), (C<sub>1</sub>-C<sub>3</sub>)alkyl(difluoromethylene)(C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)acyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, piperazinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)acylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylamino)<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>11</sup>CO(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>11</sup> is R<sup>12</sup>O or R<sup>12</sup>R<sup>13</sup>N wherein R<sup>12</sup> and R<sup>13</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl; and R<sup>14</sup>(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>14</sup> is (C<sub>1</sub>-C<sub>6</sub>)acylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)arylpiperazino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylpiperazino, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)arylpiperidyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl or (C<sub>1</sub>-C<sub>6</sub>)acylpiperidyl;

or X<sup>1</sup> and X<sup>2</sup> may be taken together to form a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, a benzo-fused (C<sub>3</sub>-C<sub>6</sub>)cydoalkyl ring or a group of the formula (J<sub>7</sub>):




wherein the carbon atom bearing the asterisk is the carbon to which X<sup>1</sup> and X<sup>2</sup> are attached, s and t are each independently 1 or 2, and W is CF<sub>2</sub>, O, S, SO<sub>2</sub> or NR<sup>15</sup>, wherein R<sup>15</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)acyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl(C=O)-;

Q is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxy)<sub>2</sub>(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxy)<sub>2</sub>(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, wherein

each of the foregoing aryl groups may be optionally substituted by fluoro, chloro, bromo, trifluoromethyl, trifluoromethoxy, difluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or perfluoro(C<sub>1</sub>-C<sub>3</sub>)alkyl; with the proviso that when either X<sup>1</sup> or X<sup>2</sup> is CH(R<sup>5</sup>)COR<sup>6</sup> wherein R<sup>5</sup> and R<sup>6</sup> are as defined above, the other of X<sup>1</sup> or X<sup>2</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or benzyl.


[0017] One preferred embodiment of the present invention relates to a method of treating a disease, disorder or condition of the peripheral or central nervous system in a mammal, comprising the administration to a mammal a therapeutically effective amount of a compound of the formula (Ia):



or a pharmaceutically acceptable salt thereof, wherein X<sup>1</sup>, X<sup>2</sup>, Q and Z are as defined above, and n is an integer from 1 to 6.


[0018] Preferred methods of the invention comprise the administration of a compound of formula (Ia) wherein n is 2. Other preferred methods of the invention comprise the administration of a compound of formula (Ia) wherein Q is 4-methoxyphenyl or 4-phenoxyphenyl, optionally substituted by fluoro, chloro, bromo, trifluoromethyl, trifluoromethoxy, difluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or perfluoro(C<sub>1</sub>-C<sub>3</sub>)alkyl. Other preferred methods of the invention comprise the administration of a compound of formula (Ia) wherein either X<sup>1</sup> or X<sup>2</sup> is not hydrogen. Other preferred methods of the invention comprise the administration of a compound of formula (Ia) wherein Z is hydroxy, Q is 4-methoxyphenyl or 4-phenoxyphenyl, optionally substituted by fluoro, chloro, bromo, trifluoromethyl, trifluoromethoxy, difluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or perfluoro(C<sub>1</sub>-C<sub>3</sub>)alkyl, and either X<sup>1</sup> or X<sup>2</sup> is not hydrogen.

[0019] Other preferred methods of the invention comprise the administration of a compound of formula (Ia) wherein Q is 4-methoxyphenyl or 4-phenoxyphenyl and X<sup>1</sup> and X<sup>2</sup> are taken together to form (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, oxacyclohexanyl, thiocyclohexanyl, indanyl or a group of the formula:



wherein the carbon bearing the asterisk is the carbon to which X<sup>1</sup> and X<sup>2</sup> are attached and R<sup>15</sup> is (C<sub>1</sub>-C<sub>6</sub>)acyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl.

[0020] A more preferred embodiment of the present invention relates to a method of treating a disease, condition or disorder of the peripheral or central nervous system in a mammal comprising the administration of a therapeutically effective amount of a compound of the formula (Ic):



or the pharmaceutically acceptable salts thereof, wherein

X<sup>1</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl;

X<sup>2</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl; or

X<sup>1</sup> and X<sup>2</sup> taken together with the carbon atom to which they are attached form a ring selected from (C<sub>5</sub>-C<sub>7</sub>) cycloalkyl, 4-tetrahydropyranyl and 4-piperidinyl;

Z is hydroxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy; and

Y is a substituent on any of the carbon atoms of the phenyl ring capable of supporting an additional bond, preferably from 1 to 2 substituents (more preferably one substituent, most preferably one substituent in the 4-position) on the phenyl ring, independently selected from hydrogen, fluoro, chloro, trifluoromethyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoromethoxy, difluoromethoxy and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

**[0021]** The more preferred methods of the invention comprise the administration of a compound of formula (Ic) wherein Y is hydrogen, fluoro or chloro, preferably 4-fluoro or 4-chloro. Other more preferred methods comprise the administration of a compound of formula (Ic) wherein X<sup>1</sup> and X<sup>2</sup> taken together with the carbon atom to which they are attached form a cyclopentyl or 4-tetrahydropyranyl ring.

**[0022]** Other preferred methods of the invention comprise the administration of a compound of formula (Ic) wherein X<sup>1</sup> and X<sup>2</sup> are both methyl. Other preferred methods comprise the administration of a compound of formula (Ic) wherein Z is hydroxy.


**[0023]** Specifically, the most preferred methods of the invention comprise the administration of a compound of formula (Ic) selected from the group consisting of:

- 3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoylcyclopentyl) amino]-propionic acid ethyl ester;
- 3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoylcyclopentyl) amino]propionic acid;
- 3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoyl-1-methylethyl) amino]propionic acid ethyl ester;
- 3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoyl-1-methylethyl) amino]propionic acid;
- 3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoyltetrahydropyran-4-yl)-amino]propionic acid;
- 3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoyltetrahydropyran-4-yl)-amino]propionic acid ethyl ester;
- 3-[[4-(4-chlorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoyltetrahydropyran-4-yl)-amino]propionic acid;
- 3-[[4-(4-chlorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoyltetrahydropyran-4-yl)-amino]propionic acid ethyl ester;
- 3-[[4-(4-hydroxycarbamoyltetrahydropyran-4-yl)-(4-phenoxybenzenesulfonyl) amino]-propionic acid;
- 3-[[4-(4-hydroxycarbamoyltetrahydropyran-4-yl)-(4-phenoxybenzenesulfonyl) amino]propionic acid ethyl ester;
- 3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoylpiperidin-4-yl)amino]propionic acid ethyl ester;
- 3-[[4-(4-chlorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoyl-1-methylethyl) amino]-propionic acid;
- 3-[[4-(4-chlorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoyl-1-methylethyl) amino]-propionic acid ethyl ester;
- 3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoylcyclohexyl) amino]-propionic acid;
- 3-[(1-hydroxycarbamoylcyclopentyl)-(4-phenoxybenzenesulfonyl) amino]propionic acid;
- 3-[[4-(4-chlorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoylcyclopentyl) amino]-propionic acid

and pharmaceutically acceptable salts thereof.


**[0024]** The methods of the invention also encompass methods of treating or preventing comprising administering a prodrug of a compound of formula (I). A compound of formula (I) having a free amino, amido, hydroxy or carboxylic acid group can be converted into a prodrug. Prodrugs include compounds wherein an amino acid residue, or a polypeptide chain of two or more (e.g., two, three or four) amino acid residues which are covalently joined through peptide bonds to free amino, hydroxy or carboxylic acid groups of compounds of formula (I). The amino acid residues include

the 20 naturally occurring amino acids commonly designated by three letter symbols and also include, 4-hydroxyproline, hydroxylysine, demosine, isodemosine, 3-methylhistidine, norvalin, beta-alanine, gamma-aminobutyric acid, citrulline, homocysteine, homoserine, ornithine and methionine sulfone. Prodrugs also include compounds wherein carbonates, carbamates, amides and alkyl esters which are covalently bonded to the above substituents of formula (I) through the carbonyl carbon prodrug sidechain. Prodrugs also include compounds of formula (I) in which the hydroxamic acid and carbonyl moiety when taken together, for example, form a group of the formula (Id):




wherein  $X^1$ ,  $X^2$  and  $Y$  are as defined above and  $U$  and  $V$  are independently carbonyl, methylene,  $SO_2$  or  $SO_3$ , and  $b$  is an integer from one to three wherein each methylene group is optionally substituted with hydroxy.

[0025] In addition to the foregoing preferred methods, further preferred embodiments of the present invention relates to a method of treatment of a condition, disease or disorder of the peripheral or central nervous system in a mammal comprising the administration of a compound of the formula (Ib):



wherein  $X^1$ ,  $X^2$  and  $Q$  are as defined above.

[0026] Other preferred methods comprise the administration of a compound of formula (Ib) wherein  $X^1$  and  $X^2$  are taken together to form a  $(C_3-C_6)$ cycloalkyl or benzo-fused  $(C_3-C_6)$ cycloalkyl ring or a group of formula  $J_7$ :




wherein the carbon atom bearing the asterisk is the carbon to which  $X^1$  and  $X^2$  are attached,  $s$  and  $t$  are independently 1 or 2;  $W$  is  $CF_2$ ,  $S$ ,  $O$  or  $NR^{16}$  and  $R^{16}$  is hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl,  $(C_2-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfonyl,  $(C_6-C_{10})$ arylsulfonyl or acyl.

[0027] Other preferred methods comprise the administration of a compound of formula (Ib) wherein  $X^1$  and  $X^2$  are taken together to form a  $(C_3-C_6)$ cycloalkyl or benzo-fused  $(C_3-C_6)$ cycloalkyl ring. Other preferred methods comprise the administration of a compound of formula (Ib) wherein  $Q$  is  $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryl $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryloxy  $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryloxy $(C_2-C_9)$ heteroaryl,  $(C_2-C_9)$ heteroaryl,  $(C_2-C_9)$ heteroaryl $(C_2-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryl  $(C_2-C_9)$ heteroaryl,  $(C_2-C_9)$ heteroaryl $(C_6-C_{10})$ aryl or  $(C_2-C_9)$ heteroaryloxy $(C_6-C_{10})$ aryl. Other preferred methods comprise the administration of a compound of formula (Ib) wherein  $Q$  is  $(C_6-C_{10})$ aryloxy $(C_6-C_{10})$ aryl. Other preferred methods comprise the administration of a compound of formula (Ib) wherein  $X^1$  and  $X^2$  are each independently  $(C_1-C_6)$ alkyl.

[0028] Other more preferred methods comprise the administration of a compound of formula (Ib) wherein  $X^1$  and  $X^2$  are taken together to form a  $(C_3-C_6)$ cycloalkyl or benzo-fused  $(C_3-C_6)$ cycloalkyl ring or a group of the formula  $(J_7)$ :





5

wherein the carbon atom bearing the asterisk is the carbon to which  $X^1$  and  $X^2$  are attached,  $s$  and  $t$  are independently 1 or 2 and  $Q^9$  is  $\text{CF}_2$ , S, O or  $\text{NR}^{16}$  wherein  $\text{R}^{16}$  is hydrogen,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkylsulfonyl}$ ,  $(\text{C}_6\text{-C}_{10})\text{arylsulfonyl}$  or acyl; and  $Q$  is  $(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryloxy}(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryloxy}(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_6\text{-C}_{10})\text{aryl}$  or  $(\text{C}_2\text{-C}_9)\text{heteroaryloxy}(\text{C}_6\text{-C}_{10})\text{aryl}$ .

**[0029]** More preferred methods comprise the administration of a compound of formula (Ib) wherein  $X^1$  and  $X^2$  are taken together to form a  $(\text{C}_3\text{-C}_6)\text{cydoalkyl}$  or benzo-fused  $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$  ring; and  $Q$  is  $(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryloxy}(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryloxy}(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_2\text{-C}_9)\text{heteroaryl}$  or  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_6\text{-C}_{10})\text{aryl}$ .

**[0030]** More preferred methods comprise the administration of a compound of formula (Ib) wherein  $X^1$  and  $X^2$  are each independently  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ; and  $Q$  is  $(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryloxy}(\text{C}_6\text{-C}_{10})\text{aryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryloxy}(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_2\text{-C}_9)\text{heteroaryl}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_6\text{-C}_{10})\text{aryl}$  or  $(\text{C}_2\text{-C}_9)\text{heteroaryloxy}(\text{C}_6\text{-C}_{10})\text{aryl}$ . More preferred methods utilize compounds of formula (Ib) wherein  $X^1$  and  $X^2$  are each independently  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ; and  $Q$  is  $(\text{C}_6\text{-C}_{10})\text{aryloxy}(\text{C}_6\text{-C}_{10})\text{aryl}$ .

**[0031]** Further, in addition to the methods stated above, other preferred methods comprise the administration of a compound of formula (Ia), *supra*, wherein  $n$  is 1 and either of  $\text{R}^1$  or  $\text{R}^2$  is hydrogen. Other preferred methods comprise the administration of a compound of formula (Ia) wherein  $Z$  is alkoxy,  $Q$  is 4-methoxyphenyl or 4-phenoxyphenyl and either  $X^1$  or  $X^2$  is not hydrogen. More preferred methods comprise the administration of a compound of formula (Ia) wherein  $n$  is 2,  $Q$  is 4-methoxyphenyl or 4-phenoxyphenyl,  $\text{R}^1$  and  $\text{R}^2$  taken together to form piperazinyl,  $(\text{C}_1\text{-C}_6)\text{alkylpiperazinyl}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl piperazinyl}$  or  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_1\text{-C}_6)\text{alkylpiperazinyl}$ , and either  $X^1$  or  $X^2$  is not hydrogen or both  $X^1$  and  $X^2$  are not hydrogen. More preferred methods comprise the administration of a compound of formula (Ia) wherein  $n$  is 2,  $Q$  is 4-methoxyphenyl or 4-phenoxyphenyl,  $\text{R}^1$  is hydrogen or  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $\text{R}^2$  is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either  $X^1$  or  $X^2$  is not hydrogen or both  $X^1$  and  $X^2$  are not hydrogen. More preferred methods comprise the administration of a compound of formula (Ia) wherein  $n$  is 1,  $Q$  is 4-methoxyphenyl or 4-phenoxyphenyl,  $\text{R}^1$  is hydrogen,  $\text{R}^2$  is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either  $X^1$  or  $X^2$  is not hydrogen or both  $X^1$  and  $X^2$  are not hydrogen. More preferred methods comprise the administration of a compound of formula (Ia) wherein  $n$  is 2,  $Q$  is 4-methoxyphenyl,  $\text{R}^1$  is hydrogen or  $(\text{C}_1\text{-C}_6)\text{alkyl}$  and  $\text{R}^2$  is  $\text{R}^3(\text{C}_2\text{-C}_6)\text{alkyl}$  wherein  $\text{R}^3$  is morpholino, thiomorpholino, piperidino, pyrrolidino,  $(\text{C}_1\text{-C}_6)\text{acylpiperazino}$ ,  $(\text{C}_1\text{-C}_6)\text{alkylpiperazino}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl-piperazino}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl-piperazino}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_1\text{-C}_6)\text{alkylpiperazino}$  or  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_1\text{-C}_6)\text{alkylpiperazino}$  and either  $X^1$  or  $X^2$  is not hydrogen or both  $X^1$  and  $X^2$  are not hydrogen. More preferred methods comprise the administration of a compound of formula (Ia) wherein  $n$  is 1,  $Q$  is 4-methoxyphenyl or 4-phenoxyphenyl,  $\text{R}^1$  is hydrogen,  $\text{R}^2$  is  $\text{R}^3(\text{C}_2\text{-C}_6)\text{alkyl}$  wherein  $\text{R}^3$  is morpholino, thiomorpholino, piperidino, pyrrolidino,  $(\text{C}_1\text{-C}_6)\text{acylpiperazino}$ ,  $(\text{C}_1\text{-C}_6)\text{alkylpiperazino}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl-piperazino}$ ,  $(\text{C}_2\text{-C}_9)\text{heteroaryl-piperazino}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryl}(\text{C}_1\text{-C}_6)\text{alkylpiperazino}$  or  $(\text{C}_2\text{-C}_9)\text{heteroaryl}(\text{C}_1\text{-C}_6)\text{alkylpiperazino}$ , and either  $X^1$  or  $X^2$  is not hydrogen or both  $X^1$  and  $X^2$  are not hydrogen.

**[0032]** In the foregoing discussion, the term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.

**[0033]** The term "alkoxy", as used herein, includes O-alkyl groups wherein "alkyl" is as defined above.

**[0034]** The term "aryl", as used herein, unless otherwise indicated, includes an organic radical derived from an aromatic hydrocarbon by removal of one hydrogen, such as phenyl or naphthyl, optionally substituted by 1 to 3 substituents selected from the group consisting of fluoro, chloro, bromo, perfluoro $(\text{C}_1\text{-C}_6)\text{alkyl}$  (including trifluoromethyl),  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryloxy}$ , perfluoro $(\text{C}_1\text{-C}_3)\text{alkoxy}$  (including trifluoromethoxy and difluoromethoxy) and  $(\text{C}_1\text{-C}_6)\text{alkyl}$ .

**[0035]** The term "heteroaryl", as used herein, unless otherwise indicated, includes an organic radical derived from an aromatic heterocyclic compound by removal of one hydrogen, such as pyridyl, furyl, pyrrolyl, thienyl, isothiazolyl, imidazolyl, benzimidazolyl, tetrazolyl, pyrazinyl, pyrimidyl, quinolyl, isoquinolyl, benzofuryl, isobenzofuryl, benzothienyl, pyrazolyl, indolyl, isoindolyl, purinyl, carbazolyl, isoxazolyl, thiazolyl, oxazolyl, benzthiazolyl or benzoxazolyl, optionally substituted by 1 to 2 substituents selected from the group consisting of fluoro, chloro, trifluoromethyl,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ ,  $(\text{C}_6\text{-C}_{10})\text{aryloxy}$ , trifluoromethoxy, difluoromethoxy and  $(\text{C}_1\text{-C}_6)\text{alkyl}$ . Preferred heteroaryl groups include pyridyl, furyl, thienyl, isothiazolyl, pyrazinyl, pyrimidyl, pyrazolyl, isoxazolyl, thiazolyl or oxazolyl. Most preferred heteroaryl groups include pyridyl, furyl or thienyl.

[0036] The term "acyl", as used herein, unless otherwise indicated, includes a radical of the general formula RCO wherein R is alkyl, alkoxy, aryl, arylalkyl or arylalkyloxy and the terms "alkyl" or "aryl" are as defined above.

[0037] The term "acyloxy", as used herein, includes O-acyl groups wherein "acyl" is defined above.

5 [0038] The term "treating" refers to, and includes, reversing, alleviating, inhibiting the progress of, or preventing a disease, disorder or condition, or one or more symptoms thereof; and "treatment" and "therapeutically" refer to the act of treating, as defined above.

[0039] A "therapeutically effective amount" is any amount of any of the compounds utilized in the course of practicing the invention provided herein that is sufficient to reverse, alleviate, inhibit the progress of, or prevent a disease, disorder or condition, or one or more symptoms thereof.

10 [0040] The methods of the invention comprise the administration of a compound of formula (I) which may have chiral centers and therefore exist in different enantiomeric forms. This invention relates to all optical isomers, tautomers and stereoisomers of the compounds of formula (I) and mixtures thereof.

[0041] The present invention also relates to a method comprising the administration of a pharmaceutically acceptable acid addition salt of a compound of the formula (I). The possible acids which are used to prepare the pharmaceutically acceptable acid addition salts of the aforementioned base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, add phosphate, acetate, lactate, citrate, acid citrate, tartrate, bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate (i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts.

20 [0042] The invention also relates to a method comprising the administration of a base addition salt of a compound of formula (I). The chemical bases that may be used as reagents to prepare pharmaceutically acceptable base salts of those compounds of formula (I) that are acidic in nature are those that form non-toxic base salts with such compounds. Such non-toxic base salts include, but are not limited to those derived from such pharmacologically acceptable cations such as alkali metal cations (e.g., potassium and sodium) and alkaline earth metal cations (e.g., calcium and magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine-(meglumine), and the lower alkanolammonium and other base salts of pharmaceutically acceptable organic amines.

25 [0043] The subject invention also relates to a method of treatment which relates to isotopically-labeled compounds, which are identical to those recited in formula (I), but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds relating to the present invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as  $^2\text{H}$ ,  $^3\text{H}$ ,  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{18}\text{O}$ ,  $^{17}\text{O}$ ,  $^{31}\text{P}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{18}\text{F}$ , and  $^{36}\text{Cl}$ , respectively. Compounds relating to the present invention, prodrugs thereof, and pharmaceutically acceptable salts of said compounds or of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labeled compounds of the present invention, for example those into which radioactive isotopes such as  $^3\text{H}$  and  $^{14}\text{C}$  are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e.,  $^3\text{H}$ , and carbon-14, i.e.,  $^{14}\text{C}$ , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e.,  $^2\text{H}$ , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances.

40 [0044] The present invention also relates to a pharmaceutical composition for the treatment of a disease, condition or disorder of the peripheral or central nervous system, wherein the disease, condition or disorder is Alzheimer's disease, stroke/cerebral ischemia, head trauma, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, migraine, cerebral amyloid angiopathy, AIDS, age-related cognitive decline; mild cognitive impairment or a prion disease.

45 [0045] The present invention also relates to a pharmaceutical composition for treating of a disease, disorder or condition, wherein the disease, condition or disorder is arthritis (including osteoarthritis and rheumatoid arthritis), inflammatory bowel disease, Crohn's disease, emphysema, chronic obstructive pulmonary disease, Alzheimer's disease, organ transplant toxicity, cachexia, allergic reactions, allergic contact hypersensitivity, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, osteoporosis, loosening of artificial joint implants, atherosclerosis (including atherosclerotic plaque rupture), aortic aneurysm (including abdominal aortic aneurysm and brain aortic aneurysm), congestive heart failure, myocardial infarction, stroke, cerebral ischemia, head trauma, spinal cord injury, neurodegenerative disorders (acute and chronic), inflammatory and autoimmune disorders, Huntington's disease, Parkinson's disease, migraine, depression, peripheral neuropathy, pain, cerebral amyloid angiopathy, nootropic or cognitive enhancement, amyotrophic lateral sclerosis, multiple sclerosis, ocular angiogenesis, corneal injury, macular degeneration, abnormal wound healing, burns, diabetes, tumor invasion, tumor growth, tumor metastasis, corneal scarring, scleritis, AIDS, sepsis, septic shock, other diseases characterized by metalloproteinase activity or other diseases characterized by mammalian repolysin activity in a mammal, including a human, comprising an amount of a compound

of formula (I) or a pharmaceutically acceptable salt thereof effective in such treatments and a pharmaceutically acceptable carrier.

[0046] This invention also encompasses pharmaceutical compositions containing a prodrug of a compound of the formula (I). This invention also encompasses methods of treating or preventing disorders that can be treated or prevented by the inhibition of matrix metalloproteinases or the inhibition of mammalian reprotolysin comprising a administering prodrug of compounds of the formula (I). A compound of formula (I) having a free amino, amido, hydroxy or carboxylic group can be converted into a prodrug. Prodrugs include compounds wherein an amino acid residue, or a polypeptide chain of two or more (e.g., two, three or four) amino acid residues which are covalently joined through peptide bonds to free amino, hydroxy or carboxylic acid groups of compounds of formula (I). The amino acid residues include the 20 naturally occurring amino acids commonly designated by three letter symbols and also include, 4-hydroxyproline, hydroxylysine, demosine, isodemossine, 3-methylhistidine, norvalin, beta-alanine, gamma-aminobutyric acid, citrulline, homocysteine, homoserine, ornithine and methionine sulfone. Prodrugs also include compounds wherein carbonates, carbamates, amides and alkyl esters which are covalently bonded to the above substituents of formula I through the carbonyl carbon prodrug side chain.

[0047] One of ordinary skill in the art will appreciate that the methods of the invention are useful in treating a diverse array of diseases. One of ordinary skill in the art will also appreciate that when using the methods of the invention in the treatment of a specific disease that the methods of the invention may be combined with various existing therapeutic methods and agents used for that disease.

[0048] The present invention also relates to combination therapies using, or combination pharmaceutical compositions comprising, a compound of formula (I) and, e.g., a standard non-steroidal anti-inflammatory drug (NSAID'S), such as piroxicam, diclofenac, a propionic acid, such as naproxen, flubiprofen, fenoprofen, ketoprofen and ibuprofen, a fenamate, such as mefenamic acid, indomethacin, sulindac, apazone, a pyrazolone, such as phenylbutazone; a salicylate, such as aspirin; an analgesic or intraarticular therapy, such as a corticosteroid and a hyaluronic acid, such as hyalgan and synvisc; an immune suppressant, such as cyclosporin, interferon, etc., e.g., in organ transplant therapy; a TNF- $\alpha$  inhibitor agent, such as an anti-TNF monoclonal antibody, a TNF receptor immunoglobulin molecule (such as Enbrel®), low dose methotrexate, lefunimide, hydroxychloroquine, d-penicillamine, auranofin, parenteral gold, oral gold, etc.

[0049] The methods of present invention also relate to combination therapies using or combination pharmaceutical compositions comprising, a compound of formula (I) and, e.g., a CNS agent or agents, such as an antidepressant (e.g., sertraline, fluoxetine, paroxetine, etc.); an anti-Parkinsonian drug, such as deprenyl, L-dopa, requip, miratex, etc.; a MAOB inhibitor, such as selegine, rasagiline, etc.; a COMT inhibitor, such as tolcapone (i.e., Tasmar); an A-2 inhibitor; a dopamine reuptake inhibitor; an NMDA antagonist; a nicotine agonist; a dopamine agonist; an inhibitor of neuronal nitric oxide synthase; an anti-Alzheimer's drug; an acetylcholinesterase inhibitor, such as metrifonate, donepezil (i.e., Aricept), Exelon (i.e., ENA 713 or rivastigmine), etc.; tetrahydroaminoacridine (i.e., Tacrine, Cognex, or THA); a COX-1 or COX-2 inhibitor, such as celecoxib (i.e., Celebrex), rofecoxib (i.e., Vioxx), etc.; propentofylline; an anti-stroke medication; an NR2B selective antagonist; a glycine site antagonist; a neutrophil inhibitory factor (NIF), etc. The methods of the present invention further relate to combination therapies using, or combination pharmaceutical compositions comprising, a compound of formula (I) and, e.g., an estrogen; a selective estrogen modulator, such as estrogen, raloxifene, tamoxifene, droloxifene, lasofoxifene, etc. The methods of the present invention also relate to combination therapies using, or combination pharmaceutical compositions comprising, a compound of formula (I) and, e.g., an agent that results in reduction of A $\beta$ 1-40/1-42, such as an amyloid aggregation inhibitor, a secretase inhibitor, etc.

[0050] Further, the methods of present invention also relate to combination therapies using, or combination pharmaceutical compositions comprising, a compound of formula (I) and, e.g., an osteoporosis agents such as droloxifene or fosomax and immunosuppressant agents such as FK-506 and rapamycin. The methods of present invention also relate to combination therapies using, or combination pharmaceutical compositions comprising, a compound of formula (I) and, e.g., an anticancer agent, such as endostatin and angiostatin; a cytotoxic drug, such as adriamycin, daunomycin, cis-platinum, etoposide, taxol, taxotere or an alkaloid, such as vincristine; an antimetabolite, such as methotrexate; a cardiovascular agent, such as calcium channel blockers; a lipid lowering agent, such as a statin, a fibrate, a beta-blocker, an ACE inhibitor, an angiotensin-2 receptor antagonist or a platelet aggregation inhibitor. The methods of present invention also relate to combination therapies comprising the administration of compounds of formula (I) and another treatment, such as, e.g., fetal implant surgery treatment, gene therapy, etc.

#### DETAILED DESCRIPTION OF THE INVENTION

[0051] Matrix metalloproteinase inhibitors, including MMP-2 and MMP-9 selective inhibitors, utilized in accordance with the methods of the present invention, can be prepared according to methods well known to those of ordinary skill in the art. Specifically, methods for the preparation of the  $\alpha$ -sulfonylamino hydroxamic acid matrix metalloproteinase inhibitors used in the methods of the present invention have been described in PCT Publication WO 96/27583, published

March 7, 1996, filed as U.S. Patent Application No. 08/401,049 on March 8, 1995; PCT Publication WO 98/33768, published August 6, 1998, filed as U.S. Patent Application No. 09/355,163 on January 12, 1998; and PCT Publication WO 99/07675, published February 18, 1999, filed as U.S. Patent Application No. 60/055,207 on August 8, 1997. The content of the foregoing U.S. patent applications are hereby incorporated in their entirety by reference. Isotopically labeled compounds of formula (I) of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples and Preparations of the foregoing incorporated applications, by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

[0052] The following additional U.S. patent applications relating specifically to other matrix metalloproteinase inhibitors with broad activity and their methods of preparation are also hereby incorporated by reference: U.S. Patent Application No. 09/154,969, filed September 17, 1998 which refers to non-selective cyclic aryl sulfonamino hydroxamic acids; U.S. Patent Application No. 08/855,023, filed May 13, 1997 which refers to cyclic imides as MMP inhibitors; U.S. Patent Application No. 08/881,092, filed July 9, 1997, which refers to cyclic sulfone hydroxamic acids useful as MMP inhibitors; U.S. Patent Application No. 09/242,504, filed July 25, 1997, which refers to aryl sulfonamino hydroxamic acids; U.S. Patent Application No. 08/892,417 filed July 14, 1997 which refers to phosphanates that are useful as MMP inhibitors; U.S. Patent Application No. 09/125,055, filed January 16, 1998, which refers to cyclic aryl sulfonamino hydroxamic acids useful in MMP inhibitors; U.S. Patent Application No. 09/341,226, filed January 27, 1998, which refers to N-hydroxy- $\beta$ -sulfonyl propionamides useful as MMP inhibitors; U.S. Patent Application No. 09/331,275, filed December 18, 1997, which refers to cyclic sulfones as useful MMP inhibitors; U.S. Patent Application Ser. No. 09/233,883, filed January 20, 1999; U.S. Patent Application No. 09/216,402, filed December 18, 1998; U.S. Patent Application No. 09/130,922, filed August 6, 1998; (U.S. Patent Application No. 09/290,022, filed April 9, 1999, U.S. Patent Application No. 09/287,930, filed April 7, 1999; and U.S. Patent Application No. 09/287,508, filed April 7, 1999.

[0053] Further applications relating to matrix metalloproteinase inhibitor compounds hereby incorporated by reference are the following: PCT Application No. PCT/EP98/08565, filed December 23, 1998; and PCT Application No. PCT/EP98/06640, filed October 9, 1998.

[0054] The ability of matrix metalloproteinase inhibitors or their pharmaceutically acceptable salts (hereinafter also referred to as the inhibitors utilized in the present invention) to inhibit matrix metalloproteinases and a demonstration of their effectiveness for treating diseases of the peripheral and central nervous system is shown by the following *in vitro* assay tests.

[0055] In studying the regulation of genes in cytokine-stimulated human astrocytes, over-expression of one of the endogenous inhibitors of MMP, TIMP-1 (Tissue Inhibitor of Metalloproteinase 1) has been detected. In AD and age-matched controlled brains, increased expression of TIMP-1, MMP-2 and MMP-9 has been observed. Increased expression of two of the MMP's known to be inhibited by TIMP-1, MMP-2 and MMP-9, has been demonstrated (See, Table 1). Brain tissue from the hippocampus and superior frontal gyrus (SFG) have been selected for analysis because they are severely affected by AD degeneration; cerebellum was studied because it is relatively unaffected by the disease process. It has been determined that the expression of MMP-2, MMP-9 and TIMP-1 are all increased in the hippocampus and SFG. MMP-2 and -9 are also up-regulated in the cerebellum but perhaps not to as great an extent.

Table 1.

Ratio of mRNA Expression in AD v. Age-matched Control Brain Regions.			
	Hippocampus Region	SFG Region	Cerebellum Region
MMP-2	2.5 (p=0.002)	2.4 (p=0.006)	1.5 (p=0.033)
MMP-9	3.3 (p=0.040)	6.4 (p=0.006)	2.5 (p=0.045)
TIMP-1	1.8 (p=0.006)	2.2 (p=0.003)	1.1 (p=0.541)

#### Biological Assay

[0056] The following assays may be used to identify matrix metalloproteinase inhibitors.

#### Inhibition of Human Collagenase (MMP-1)

[0057] Human recombinant collagenase is activated with trypsin using the following ratio: 10  $\mu$ g trypsin per 100  $\mu$ g of collagenase. The trypsin and collagenase are incubated at room temperature for 10 minutes then a five fold excess (50  $\mu$ g/10  $\mu$ g trypsin) of soybean trypsin inhibitor is added.

[0058] 10 mM stock solutions of inhibitors are made up in dimethyl sulfoxide and then diluted using the following Scheme:

10 mM → 120 μM → 12 μM → 1.2 μM → 0.12 μM

[0059] Twenty-five microliters of each concentration is then added in triplicate to appropriate wells of a 96 well microfluor plate. The final concentration of inhibitor will be a 1:4 dilution after addition of enzyme and substrate. Positive controls (enzyme, no inhibitor) are set up in wells D1-D6 and blanks (no enzyme, no inhibitors) are set in wells D7-D12.

[0060] Collagenase is diluted to 240 ng/ml and 25 μl is then added to appropriate wells of the microfluor plate. Final concentration of collagenase in the assay is 60 ng/ml.

[0061] Substrate (DNP-Pro-Cha-Gly-Cys(Me)-His-Ala-Lys(NMA)-NH<sub>2</sub>) is made as a 5 mM stock in dimethyl sulfoxide and then diluted to 20 μM in assay buffer. The assay is initiated by the addition of 50 μl substrate per well of the microfluor plate to give a final concentration of 10 μM.

[0062] Fluorescence readings (360 nm excitation, 460 nm emission) were taken at time 0 and then at 20 minute intervals. The assay is conducted at room temperature with a typical assay time of 3 hours.

[0063] Fluorescence vs time is then plotted for both the blank and collagenase containing samples (data from triplicate determinations is averaged). A time point that provides a good signal (the blank) and that is on a linear part of the curve (usually around 120 minutes) is chosen to determine IC<sub>50</sub> values. The zero time is used as a blank for each compound at each concentration and these values are subtracted from the 120 minute data. Data is plotted as inhibitor concentration vs % control (inhibitor fluorescence divided by fluorescence of collagenase alone x 100). IC<sub>50</sub>'s are determined from the concentration of inhibitor that gives a signal that is 50% of the control.

[0064] If IC<sub>50</sub>'s are reported to be <0.03 μM then the inhibitors are assayed at concentrations of 0.03 μM, 0.003 μM, 0.0003 μM and 0.00003 μM.

#### Inhibition of Gelatinase (MMP-2)

[0065] Inhibition of gelatinase activity is assayed using the MCA-Pro-Leu-Gly-Leu-Dpa-Ala-Arg-NH<sub>2</sub> substrate (10 μM) under the same conditions as inhibition of human collagenase (MMP-1).

[0066] 72kD gelatinase is activated with 1 mM APMA (p-aminophenyl mercuric acetate) for 16-18 hours at 4°C and is diluted to give a final concentration in the assay of 25 ng/ml. Inhibitors are diluted as for inhibition of human collagenase (MMP-1) to give final concentrations in the assay of 30 μM, 3 μM, 0.3 μM and 0.03 μM. Each concentration is done in triplicate.

[0067] Fluorescence readings (320 nm excitation, 390 emission) are taken at time zero and then at 15 minutes intervals for 3 hours.

[0068] IC<sub>50</sub>'s are determined as per inhibition of human collagenase (MMP-1). If IC<sub>50</sub>'s are reported to be less than 0.03 μM, then the inhibitors are assayed at final concentrations of 0.03 μM, 0.003 μM, 0.0003 μM and 0.00003 μM.

#### Inhibition of Stromelysin Activity (MMP-3)

[0069] Inhibition of stromelysin activity is assayed using Mca-Arg-Pro-Lys-Pro-Val-Glu-Nva-Trp-Arg-Lys(Dnp)-NH<sub>2</sub> (3 μM) under conditions similar as in inhibition of human collagenase (MMP-1).

[0070] Human stromelysin is activated for 20-24 hours at 37°C with 2 mM APMA (p-aminophenyl mercuric acetate) and is diluted to give a final concentration in the assay of 50 ng/ml. Inhibitors are diluted as for inhibition of human collagenase (MMP-1) to give final concentrations in the assay of 30 μM, 3 μM, 0.3 μM, and 0.03 μM. Each concentration is done in triplicate.

[0071] Fluorescence readings (320 nm excitation, 390 emission) are taken at time zero and then at 15 minute intervals for 3 hours.

[0072] IC<sub>50</sub>'s are determined as per inhibition of human collagenase (MMP-1). If IC<sub>50</sub>'s are reported to be less than 0.03 μM, then the inhibitors are assayed at final concentrations of 0.03 μM, 0.003 μM, 0.0003 μM, and 0.00003 μM.

#### Inhibition of Gelatinase (MMP-9)

[0073] Inhibition of gelatinase activity is assayed using the MCA-Pro-Leu-Gly-Leu-Dpa-Ala-Arg-NH<sub>2</sub> substrate (10 μM) under the same conditions as inhibition of human collagenase (MMP-1).

[0074] 92kD gelatinase is activated with 1.5 mM APMA (p-aminophenyl mercuric acetate) for 2 hours at 37°C and is diluted to give a final concentration in the assay of 25 ng/ml. Inhibitors are diluted as for inhibition of human collagenase (MMP-1) to give final concentrations in the assay of 30 μM, 3 μM, 0.3 μM and 0.03 μM. Each concentration is done in triplicate.

[0075] Fluorescence readings (320 nm excitation, 390 emission) are taken at time zero and then at 15 minutes intervals for 3 hours.

[0076]  $IC_{50}$ 's are determined as per inhibition of human collagenase (MMP-1). If  $IC_{50}$ 's are reported to be less than 0.03  $\mu$ M, then the inhibitors are assayed at final concentrations of 0.03  $\mu$ M, 0.003  $\mu$ M, 0.0003  $\mu$ M and 0.00003  $\mu$ M.

#### Inhibition of MMP-13

[0077] Human recombinant MMP-13 is activated with 2 mM APMA (p-aminophenyl mercuric acetate) for 1.5 hours, at 37°C and is diluted to 240  $\mu$ g/ml in assay buffer (50 mM Tris, pH 7.5, 200 mM sodium chloride, 5 mM calcium chloride, 20  $\mu$ M zinc chloride, 0.02% brij). Twenty-five microliters of diluted enzyme is added per well of a 96 well microfluor plate. The enzyme is then diluted in a 1:4 ratio in the assay by the addition of inhibitor and substrate to give a final concentration in the assay of 60  $\mu$ g/ml.

[0078] 10 mM stock solutions of inhibitors are made up in dimethyl sulfoxide and then diluted in assay buffer as per the inhibitor dilution scheme for inhibition of human collagenase (MMP-1): Twenty-five microliters of each concentration is added in triplicate to the microfluor plate. The final concentrations in the assay are 30  $\mu$ M, 3  $\mu$ M, 0.3  $\mu$ M, and 0.03  $\mu$ M.

[0079] Substrate (Dnp-Pro-Gly-Cys(Me)-His-Ala-Lys(NMA)-NH<sub>2</sub>) is prepared as for inhibition of human collagenase (MMP-1) and 50  $\mu$ l is added to each well to give a final assay concentration of 10  $\mu$ M. Fluorescence readings (360 nm excitation; 450 emission) are taken at time 0 and every 5 minutes for 1 hour.

[0080] Positive controls consist of enzyme and substrate with no inhibitor and blanks consist of substrate only.

[0081]  $IC_{50}$ 's are determined as per inhibition of human collagenase (MMP-1). If  $IC_{50}$ 's are reported to be less than 0.03  $\mu$ M, inhibitors are then assayed at final concentrations of 0.03  $\mu$ M, 0.003  $\mu$ M, 0.0003  $\mu$ M and 0.00003  $\mu$ M.

#### Collagen film MMP-13 Assay

[0082] Rat type I collagen is radiolabeled with <sup>14</sup>C acetic anhydride (T.E. Cawston and A.J. Barrett, *Anal. Biochem.*, 99, 340-345 (1979)) and used to prepare 96 well plates containing radiolabeled collagen films (Barbara Johnson-Wint, *Anal. Biochem.*, 104, 175-181 (1980)). When a solution containing collagenase is added to the well, the enzyme cleaves the insoluble collagen which unwinds and is thus solubilized. Collagenase activity is directly proportional to the amount of collagen solubilized, determined by the proportion of radioactivity released into the supernatant as measured in a standard scintillation counter. Collagenase inhibitors are, therefore, compounds which reduce the radioactive counts released with respect to the controls with no inhibitor present. One specific embodiment of this assay is described in detail below.

[0083] For determining the selectivity of compounds for MMP-13 versus MMP-1 using collagen as a substrate, the following procedure is used. Recombinant human proMMP-13 or proMMP-1 is activated according to the procedures outlined above. The activated MMP-13 or MMP-1 is diluted to 0.6  $\mu$ g/ml with buffer (50 mM Tris pH 7.5, 150 mM NaCl, 10 mM CaCl<sub>2</sub>, 1  $\mu$ M ZnCl<sub>2</sub>, 0.05% Brij-35, 0.02% sodium azide).

[0084] Stock solutions of test compound (10mM) in dimethylsulfoxide are prepared. Dilutions of the test compounds in the Tris buffer, above, are made to 0.2, 2.0, 20, 200, 2000 and 20000 nM.

[0085] 100  $\mu$ l of appropriate drug dilution and 100  $\mu$ l of diluted enzyme are pipetted into wells of a 96 well plate containing collagen films labeled with <sup>14</sup>C-collagen. The final enzyme concentration is 0.3  $\mu$ g/ml while the final drug concentration is 0.1, 1.0, 10, 100, 1000 nM. Each drug concentration and control is analyzed in triplicate. Triplicate controls are also run for the conditions in which no enzyme is present and for enzyme in the absence of any compound. [0086] The plates are incubated at 37°C for a time period such that around 30 - 50% of the available collagen is solubilized - determined by counting additional control wells at various time points. In most cases around 9 hours of incubation are required. When the assay has progressed sufficiently, the supernatant from each well is removed and counted in a scintillation counter. The background counts (determined by the counts in the wells with no enzyme) are subtracted from each sample and the % release calculated in relation to the wells with enzyme only and no inhibitor. The triplicate values for each point are averaged and the data graphed as percent release versus drug concentration.  $IC_{50}$ 's are determined from the point at which 50% inhibition of release of radiolabeled collagen is obtained.

[0087] To determine the identity of the active collagenases in cartilage conditioned medium, assays were carried out using collagen as a substrate, cartilage conditioned medium containing collagenase activity and inhibitors of varying selectivity. The cartilage conditioned medium was collected during the time at which collagen degradation was occurring and thus is representative of the collagenases responsible for the collagen breakdown. Assays were carried out as outlined above except that instead of using recombinant MMP-13 or recombinant MMP-1, cartilage conditioned medium was the enzyme source.

#### IL-1 Induced Cartilage Collagen Degradation From Bovine Nasal Cartilage

[0088] This assay uses bovine nasal cartilage explants which are commonly used to test the efficacy of various compounds to inhibit either IL-1 induced proteoglycan degradation or IL-1 induced collagen degradation. Bovine nasal

cartilage is a tissue that is very similar to articular cartilage, i.e. chondrocytes surrounded by a matrix that is primarily type II collagen and aggrecan. The tissue is used because it: (1) is very similar to articular cartilage, (2) is readily available, (3) is relatively homogeneous, and (4) degrades with predictable kinetics after IL-1 stimulation.

[0089] Two variations of this assay have been used to assay compounds. Both variations give similar data. The two variations are described below:

#### Variation 1

[0090] Three plugs of bovine nasal cartilage (approximately 2 mm diameter x 1.5 mm long) are placed into each well of a 24 well tissue culture plate. One ml of serumless medium is then added to each well. Compounds are prepared as 10 mM stock solutions in DMSO and then diluted appropriately in serumless medium to final concentrations, e.g., 50, 500 and 5000 nM. Each concentration is assayed in triplicate.

[0091] Human recombinant IL-1 $\alpha$  (5ng/mL) (IL-1) is added to triplicate control wells and to each well containing drug. Triplicate control wells are also set up in which neither drug nor IL-1 are added. The medium is removed and fresh medium containing IL-1 and the appropriate drug concentrations is added on days 6, 12, 18 and 24 or every 3 - 4 days if necessary. The media removed at each time point is stored at -20°C for later analysis. When the cartilage in the IL-1 alone wells has almost completely resorbed (about day 21), the experiment is terminated. The medium is removed and stored. Aliquots (100  $\mu$ l) from each well at each time point are pooled, digested with papain and then analyzed for hydroxyproline content. Background hydroxyproline (average of wells with no IL-1 and no drug) is subtracted from each data point and the average calculated for each triplicate. The data is then expressed as a percent of the IL-1 alone average value and plotted. The IC<sub>50</sub> is determined from this plot.

#### Variation 2

[0092] The experimental set-up is the same as outlined above in Variation 1, until day 12. On day 12, the conditioned medium from each well is removed and frozen. Then one ml of phosphate buffered saline (PBS) containing 0.5  $\mu$ g/ml trypsin is added to each well and incubation continued for a further 48 hours at 37°C. After 48 hours incubation in trypsin, the PBS solution is removed. Aliquots (50  $\mu$ l) of the PBS/trypsin solution and the previous two time points (days 6 and 12) are pooled, hydrolyzed and hydroxyproline content determined. Background hydroxyproline (average of wells with no IL-1 and no drug) is subtracted from each data point and the average calculated for each triplicate. The data is then expressed as a percent of the IL-1 alone average value and plotted. The IC<sub>50</sub> is determined from this plot. In this variation, the time course of the experiment is shortened considerably. The addition of trypsin for 48 hours after 12 days of IL-1 stimulation likely releases any type II collagen that has been damaged by collagenase activity but not yet released from the cartilage matrix. In the absence of IL-1 stimulation, trypsin treatment produces only low background levels of collagen degradation in the cartilage explants.

#### Inhibition of Human 92 kD Gelatinase (MMP-9)

[0093] Inhibition of 92 kD gelatinase (MMP-9) activity is assayed using the Mca-Pro-Leu-Gly-Leu-Dpa-Ala-Arg-NH<sub>2</sub> substrate (10  $\mu$ M) under similar conditions as described above for the inhibition of human collagenase (MMP-1).

[0094] Human recombinant 92 kD gelatinase (MMP-9, gelatinase B) is activated for 2 hours with 1mM p-aminophenylmercuric acetate (from a freshly prepared 100 mM stock in 0.2 N NaOH) at 37°C.

[0095] 10 mM dimethylsulfoxide stock solutions of inhibitors are diluted serially in assay buffer (50 mM TRIS, pH 7.5, 200 mM NaCl, 5 mM CaCl<sub>2</sub>, 20  $\mu$ M ZnCl<sub>2</sub>, 0.02% BRIJ-35 (vol./vol.)) using the following scheme:

$$10 \text{ mM} \rightarrow 120 \text{ } \mu\text{M} \rightarrow 12 \text{ } \mu\text{M} \rightarrow 1.2 \text{ } \mu\text{M} \rightarrow 0.12 \text{ } \mu\text{M}$$

[0096] Further dilutions are made as necessary following this same scheme. A minimum of four inhibitor concentrations for each compound are performed in each assay. 25  $\mu$ L of each concentration is then added to triplicate wells of a black 96 well U-bottomed microfluor plate. As the final assay volume is 100  $\mu$ L, final concentrations of inhibitor are the result of a further 1:4 dilution (i.e. 30  $\mu$ M  $\rightarrow$  3  $\mu$ M  $\rightarrow$  0.3  $\mu$ M  $\rightarrow$  0.03  $\mu$ M, etc.). A blank (no enzyme, no inhibitor) and a positive enzyme control (with enzyme, no inhibitor) are also prepared in triplicate.

[0097] Activated enzyme is diluted to 100 ng/mL in assay buffer, 25  $\mu$ L per well is added to appropriate wells of the microplate. Final enzyme concentration in the assay is 25 ng/mL (0.27 nM).

[0098] A five mM dimethylsulfoxide stock solution of substrate (Mca-Pro-Leu-Gly-Leu-Dpa-Ala-Arg-NH<sub>2</sub>) is diluted in assay buffer to 20  $\mu$ M. The assay is initiated by addition of 50  $\mu$ L of diluted substrate yielding a final assay concentration of 10  $\mu$ M substrate. A zero time fluorescence reading (320 excitation; 390 emission) is immediately taken and

subsequent readings are taken every fifteen minutes at room temperature with a PerSeptive Biosystems CytoFluor Multi-Well Plate Reader with the gain at 90 units.

[0099] The average value of fluorescence of the enzyme and blank are plotted versus time. An early time point on the linear part of this curve is chosen for IC<sub>50</sub> determinations. The zero time point for each compound at each dilution is subtracted from the latter time point and the data then expressed as percent of enzyme control (inhibitor fluorescence divided by fluorescence of positive enzyme control x 100). Data is plotted as inhibitor concentration versus percent of enzyme control. IC<sub>50</sub>'s are defined as the concentration of inhibitor that gives a signal that is 50% of the positive enzyme control.

#### Aggrecanase Assay

[0100] Primary porcine chondrocytes from articular joint cartilage are isolated by sequential trypsin and collagenase digestion followed by collagenase digestion overnight and are plated at  $2 \times 10^5$  cells per well into 48 well plates with 5  $\mu$ Ci / ml 35 S (1000 Ci/mmol) sulphur in type I collagen coated plates. Cells are allowed to incorporate label into their proteoglycan matrix (approximately 1 week) at 37°C, under an atmosphere of 5% CO<sub>2</sub>.

[0101] The night before initiating the assay, chondrocyte monolayers are washed two times in DMEM/ 1% PSF/G and then allowed to incubate in fresh DMEM /1% FBS overnight.

[0102] The following morning chondrocytes are washed once in DMEM/1%PSF/G. The final wash is allowed to sit on the plates in the incubator while making dilutions.

[0103] Media and dilutions can be made as described in the Table below.

Control Media	DMEM alone (control media)
IL-1 Media	DMEM + IL-1 (5 ng/ml)
Drug Dilutions	<p>Make all compounds stocks at 10 mM in DMSO.</p> <p>Make a 100 <math>\mu</math>M stock of each compound in DMEM in 96 well plate. Store in freezer overnight.</p> <p>The next day perform serial dilutions in DMEM with IL-1 to 5 <math>\mu</math>M, 500 nM, and 50 nM.</p> <p>Aspirate final wash from wells and add 50 <math>\mu</math>l of compound from above dilutions to 450 <math>\mu</math>l of IL-1 media in appropriate wells of the 48 well plates.</p> <p>Final compound concentrations equal 500 nM, 50 nM, and 5 nM. All samples completed in triplicate with Control and IL-1 alone samples on each plate.</p>

[0104] Plates are labeled and only the interior 24 wells of the plate are used. On one of the plates, several columns are designated as IL-1 (no drug) and Control (no IL-1, no drug). These control columns are periodically counted to monitor 35S-proteoglycan release. Control and IL-1 media are added to wells (450  $\mu$ l) followed by compound (50  $\mu$ l) so as to initiate the assay. Plates are incubated at 37°C, with a 5% CO<sub>2</sub> atmosphere.

[0105] At 40-50 % release (when CPM from IL-1 media is 4-5 times control media) as assessed by liquid scintillation counting (LSC) of media samples, the assay is terminated (9-12 hours). Media is removed from all wells and placed in scintillation tubes. Scintillate is added and radioactive counts are acquired (LSC). To solubilize cell layers, 500  $\mu$ l of papain digestion buffer (0.2 M Tris, pH 7.0, 5 mM EDTA, 5 mM DTT, and 1 mg/ml papain) is added to each well. Plates with digestion solution are incubated at 60°C overnight. The cell layer is removed from the plates the next day and placed in scintillation tubes. Scintillate is then added, and samples counted (LSC).

[0106] The percent of released counts from the total present in each well is determined. Averages of the triplicates are made with control background subtracted from each well. The percent of compound inhibition is based on IL-1 samples as 0% inhibition (100% of total counts).

#### Inhibition of soluble TNF Production

[0107] The ability of the compounds or the pharmaceutically acceptable salts thereof to inhibit the cellular production/ release of TNF and, consequently, demonstrate their effectiveness for treating diseases involving the dysregulated of TNF is shown by the following *in vitro* assay:

#### Method for the evaluation of recombinant TNF $\alpha$ Converting Enzyme Activity. Preparation of recombinant TACE:

[0108] A DNA fragment coding for the signal sequence, prodomain and catalytic domain of TACE (amino acids 1-473), was amplified by polymerase chain reaction using a human lung cDNA library as a template. The amplified fragment was cloned into pFastBac vector. The DNA sequence of the insert was confirmed for both the strands. A bacmid



prepared using pFastBac in *E. coli* DH10Bac was transfected into SF9 insect cells. The virus particles were amplified to P1, P2, P3 stages. The P3 virus was infected into both Sf9 and High Five insect cells and grown at 27°C for 48 hours. The medium was collected and used for assays and further purification.

5 Preparation of fluorescent quenched substrate:

[0109] A model peptidic TNF- $\alpha$  substrate (LY-LeucineAlanineGlutamineAlanine-ValineArginineSerineSerineLysine (CMTR)-Arginine (LY=Lucifer Yellow; CMTR= 5-carboxytetramethyl Rhodamine)) was prepared and the concentration estimated by absorbance at 560 nm (E560, 60,000 M<sup>-1</sup>CM<sup>-1</sup>) according to the method of Geoghegan, KF, "Improved method for converting an unmodified peptide to an energy-transfer substrate for a proteinase." Bioconjugate Chem. 7, 385-391 (1995). This peptide encompasses the cleavage site on pro-TNF which is cleaved in vivo by TACE.

Enzyme reaction.

15 [0110] The reaction, carried out in a 96 well plate (Dynatech), was comprised of 70  $\mu$ l of buffer solution (25 mM Hepes-HCl, pH7.5, plus 20  $\mu$ M ZnCl<sub>2</sub>), 10  $\mu$ l of 100  $\mu$ M fluorescent quenched substrate, 10  $\mu$ l of a DMSO (5%) solution of test compound, and an amount of r-TACE enzyme which will cause 50% cleavage in 60 minutes - in a total volume of 100  $\mu$ l. The specificity of the enzyme cleavage at the amide bond between alanine and valine was verified by HPLC and mass spectrometry. Initial rates of cleavage were monitored by measuring the rate of increase in fluorescence at 530 nm (excitation at 409 nm) over 30 minutes. The experiment was controlled as follows: 1) for background fluorescence of substrate; 2) for fluorescence of fully cleaved substrate; 3) for fluorescence quenching or augmentation from solutions containing test compound.

20 [0111] Data was analyzed as follows. The rates from the non-test compound containing "control" reactions were averaged to establish the 100% value. The rate of reaction in the presence of test compound was compared to that in the absence of compound, and tabulated as "percent of non-test compound containing control. The results were plotted as "% of control" vs. the log of compound concentration and a half-maximal point or IC<sub>50</sub> value determined. The IC<sub>50</sub> for the above assay is a measure of the inhibition of the TNF- $\alpha$  proteolytic activity of TACE. Blockage of binding of TNF- $\alpha$  to TACE as used herein is as described in United States Patents 5,830,742, issued November 3, 1998.

30 Monocyte Assay

[0112] Human mononuclear cells were isolated from anti-coagulated human blood using a one-step Ficoll-hypaque separation technique. (2) The mononuclear cells were washed three times in Hanks balanced salt solution (HBSS) with divalent cations and resuspended to a density of 2 x 10<sup>6</sup> /ml in HBSS containing 1% BSA. Differential counts determined using the Abbott Cell Dyn 3500 analyzer indicated that monocytes ranged from 17 to 24% of the total cells in these preparations.

35 [0113] 180m of the cell suspension was aliquoted into flat bottom 96 well plates (Costar). Additions of compounds and LPS (100 ng/ml final concentration) gave a final volume of 200  $\mu$ l. All conditions were performed in triplicate. After a four hour incubation at 37°C in an humidified CO<sub>2</sub> incubator, plates were removed and centrifuged (10 minutes at approximately 250 x g) and the supernatants removed and assayed for TNF- $\alpha$  using the R&D ELISA Kit.

40 [0114] A group of preferred compounds, more preferably compounds of the formula (I), that can be identified by the methods of the present invention include those inhibitors that possess potent activity against MMP-2 and MMP-9 (preferably an IC<sub>50</sub> of less than 500nM, more preferably 100nM, most preferably 50nM) preferably wherein said MMP-2 and MMP-9 inhibitory activity is selective activity for MMP-2 and MMP-9. The compounds of formula (I) possess surprisingly selective activity against MMP-2 and MMP-9. Specifically, compounds of formula (I) have IC<sub>50</sub>'s of less than 500nM against either or both of MMP-2 and MMP-9.

45 [0115] For administration to mammals, including humans, in accordance with the methods of treatment of the present invention, for the treatment of a disorder, condition or disease of the peripheral or central nervous system, a variety of conventional routes may be used including oral, parenteral (e.g., intravenous, intramuscular or subcutaneous), buccal, anal and topical. In general, the compounds of the invention (hereinafter also known as the active compounds) will be administered at dosages between about 0.1 and 25 mg/kg body weight of the subject to be treated per day, preferably from about 0.3 to 5 mg/kg. Preferably the active compound will be administered orally or parenterally. However, some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

50 [0116] The compounds of the present invention can be administered in a wide variety of different dosage forms, in general, the therapeutically effective compounds of this invention are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

[0117] For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate,

calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof. In the case of animals, they are advantageously contained in an animal feed or drinking water in a concentration of 5-5000 ppm, preferably 25 to 500 ppm.


[0118] For parenteral administration (intramuscular, intraperitoneal, subcutaneous and intravenous use) in accordance with the present invention, a sterile injectable solution of the active ingredient is usually prepared. Solutions of a therapeutic compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably adjusted and buffered, preferably at a pH of greater than 8, if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art. In the case of animals, compounds can be administered intramuscularly or subcutaneously at dosage levels of about 0.1 to 50 mg/kg/day, advantageously 0.2 to 10 mg/kg/day given in a single dose or up to 3 divided doses.

[0119] For the methods of the present invention, the active compounds herein disclosed may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

[0120] For intranasal administration or administration by inhalation, the active compounds of the invention are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped by the patient or as an aerosol spray presentation from a pressurized container or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

## Claims

1. The use of a compound of formula (I):







or the pharmaceutically acceptable salts thereof, wherein


A is H or  $-(CH_2)_n-(C=O)-Z$ ; where n is 1 to 6; and Z is hydroxy,  $(C_1-C_6)$ alkoxy or  $NR^1R^2$  wherein  $R^1$  and  $R^2$  are each independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl, piperidyl,  $(C_1-C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ aryl piperidyl,  $(C_2-C_9)$ heteroaryl piperidyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl piperidyl,  $(C_2-C_9)$ heteroaryl  $(C_1-C_6)$ alkyl piperidyl,  $(C_1-C_6)$ acyl piperidyl,  $(C_6-C_{10})$ aryl,  $(C_2-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_2-C_9)$ heteroaryl  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl  $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryl  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl  $(C_1-C_6)$ alkyl,  $R^3(C_2-C_6)$ alkyl,  $(C_1-C_3)$ alkyl  $(CHR^3)(C_1-C_6)$ alkyl wherein  $R^3$  is hydroxy,  $(C_1-C_6)$ acyloxy,  $(C_1-C_6)$ alkoxy, piperazino,  $(C_1-C_6)$ acylamino,  $(C_1-C_6)$ alkylthio,  $(C_6-C_{10})$ arylthio,  $(C_1-C_6)$ alkylsulfinyl,  $(C_6-C_{10})$ arylsulfinyl,  $(C_1-C_6)$ alkylsulfoxyl,  $(C_6-C_{10})$ arylsulfoxyl, amino,  $(C_1-C_6)$ alkylamino,  $((C_1-C_6)alkyl)_2$  amino,  $(C_1-C_6)$ acylpiperazino,  $(C_1-C_6)$ alkylpiperazino,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkylpiperazino,  $(C_2-C_9)$ heteroaryl  $(C_1-C_6)$ alkylpiperazino, morpholino, thiomorpholino, piperidino or pyrrolidino;  $R^4(C_1-C_6)$ alkyl,  $(C_1-C_5)$ alkyl  $(CHR^4)(C_1-C_6)$ alkyl wherein  $R^4$  is piperidyl,  $(C_1-C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ aryl piperidyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkylpiperidyl,  $(C_2-C_9)$ heteroaryl piperidyl or  $(C_2-C_9)$ heteroaryl  $(C_1-C_6)$ alkylpiperidyl; and  $CH(R^5)COR^6$  wherein  $R^5$  is hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$

aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylamino)<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>7</sup>R<sup>8</sup>NCO(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sup>7</sup>OCO(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>7</sup> and R<sup>8</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl and (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl; and R<sup>6</sup> is or R<sup>9</sup>R<sup>10</sup>N wherein R<sup>9</sup> and R<sup>10</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl and (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

or R<sup>1</sup> and R<sup>2</sup>, or R<sup>7</sup> and R<sup>8</sup>, or R<sup>9</sup> and R<sup>10</sup> may be taken together to form an azetidiny, pyrrolidiny, morpholinyl, thiomorpholinyl, indolinyl, isoindolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, (C<sub>1</sub>-C<sub>6</sub>)acylpiperazinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazinyl, (C<sub>6</sub>-C<sub>10</sub>)arylpiperazinyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylpiperazinyl or a bridged diazabicycloalkyl ring selected from the group consisting of:



and



wherein

p is 1, 2 or 3;

q is 1 or 2;


r is 0 or 1;

L is hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)acyl;

X<sup>1</sup> and X<sup>2</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoromethyl, trifluoromethyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl (difluoromethylene), (C<sub>1</sub>-C<sub>3</sub>)alkyl(difluoromethylene)(C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)acyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, piperazinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)acylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylamino)<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>11</sup>CO(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>11</sup> is R<sup>12</sup>O or R<sup>12</sup>R<sup>13</sup>N wherein R<sup>12</sup> and R<sup>13</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl; and R<sup>14</sup>(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>14</sup> is (C<sub>1</sub>-C<sub>6</sub>)acylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)arylpiperazino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylpiperazino, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)arylpiperidyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylpiperi-

dyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl or (C<sub>1</sub>-C<sub>6</sub>)acylpiperidyl;

or X<sup>1</sup> and X<sup>2</sup> may be taken together to form a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, a benzo-fused (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl ring or a group of the formula (J<sub>7</sub>):




wherein the carbon atom bearing the asterisk is the carbon to which X<sup>1</sup> and X<sup>2</sup> are attached, s and t are independently 1 or 2, and W is CF<sub>2</sub>, O, S, SO<sub>2</sub> or NR<sup>15</sup>, wherein R<sup>15</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)acyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl(C=O)-;

Q is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxy)<sub>2</sub>(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxy)<sub>2</sub>(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, wherein each of the foregoing aryl groups may be optionally substituted by fluoro, chloro, bromo, trifluoromethyl, trifluoromethoxy, difluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or perfluoro(C<sub>1</sub>-C<sub>3</sub>)alkyl;

with the proviso that when either X<sup>1</sup> or X<sup>2</sup> is CH(R<sup>5</sup>)COR<sup>6</sup> wherein R<sup>5</sup> and R<sup>6</sup> are as defined above, the other of X<sup>1</sup> or X<sup>2</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or benzyl;


in the manufacture of a medicament for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to Alzheimer's disease, stroke/cerebral ischemia, head trauma, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, migraine, cerebral amyloid angiopathy, AIDS, age-related cognitive decline, mild cognitive impairment and prion diseases.

2. Use according to Claim 1, wherein A is -(CH<sub>2</sub>)<sub>n</sub>-(C=O)-Z and Z and n are as defined in Claim 1.
3. Use according to Claim 2, wherein n is 2.
4. Use according to Claim 2, wherein Q is 4-methoxyphenyl or 4-phenoxyphenyl, optionally substituted by fluoro, chloro, bromo, trifluoromethyl, trifluoromethoxy, difluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or perfluoro(C<sub>1</sub>-C<sub>3</sub>)alkyl.
5. Use according to Claim 2, wherein Z is hydroxy, Q is 4-methoxyphenyl or 4-phenoxyphenyl, optionally substituted by fluoro, chloro, bromo, trifluoromethyl, trifluoromethoxy, difluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or perfluoro(C<sub>1</sub>-C<sub>3</sub>)alkyl, and either X<sup>1</sup> or X<sup>2</sup> is not hydrogen.
6. Use according to Claim 2, wherein Q is 4-methoxyphenyl or 4-phenoxyphenyl, optionally substituted by fluoro, chloro, bromo, trifluoromethyl, trifluoromethoxy, difluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or perfluoro(C<sub>1</sub>-C<sub>3</sub>)alkyl, and X<sup>1</sup> and X<sup>2</sup> are taken together to form (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, oxacyclohexanyl, thiocyclohexanyl, indanyl or a group of the formula



wherein R<sup>15</sup> is (C<sub>1</sub>-C<sub>6</sub>)acyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl.

7. Use according to Claim 1, wherein the compound of formula (I) is a compound of formula (Ic):



or the pharmaceutically acceptable salts thereof, wherein

X<sup>1</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl;

X<sup>2</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl; or

X<sup>1</sup> and X<sup>2</sup> taken together with the carbon atom to which they are attached form a ring selected from (C<sub>5</sub>-C<sub>7</sub>) cycloalkyl 4-tetrahydropyranyl and 4-piperidiny;

Z is hydroxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy; and

Y is a substituent on any of the carbon atoms of the phenyl ring capable of supporting an additional bond, preferably from 1 to 2 substituents (more preferably one substituent, most preferably one substituent in the 4-position) on the phenyl ring, independently selected from hydrogen, fluoro, chloro, trifluoromethyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoromethoxy, difluoromethoxy and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

8. Use according to Claim 7, wherein Y is fluoro or chloro.

9. Use according to Claim 7, wherein Y is fluoro or chloro at the 4-position of the phenoxy ring.

10. Use according to Claim 7, wherein X<sup>1</sup> and X<sup>2</sup> taken together with the carbon atom to which they are attached form a cyclopentyl or 4-tetrahydropyranyl ring.

11. Use according to Claim 7, wherein Z is hydroxy.

12. Use according to Claim 7, wherein the compound of formula (Ic) is selected from the group consisting of:

3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoylcyclopentyl)amino]-propionic acid ethyl ester;

3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoylcyclopentyl) amino]propionic acid;

3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoyl-1-methylethyl) amino]propionic acid ethyl ester;

3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(1-hydroxycarbamoyl-1-methylethyl) amino]propionic acid;

3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoyltetrahydropyran-4-yl)-amino]propionic acid;

3-[[4-(4-fluorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoyltetrahydropyran-4-yl)-amino]propionic acid ethyl ester;


3-[[4-(4-chlorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoyltetrahydropyran-4-yl)-amino]propionic acid;

3-[[4-(4-chlorophenoxy)benzenesulfonyl]-(4-hydroxycarbamoyltetrahydropyran-4-yl)-amino]propionic acid ethyl ester;

- 3-[(4-hydroxycarbamoyltetrahydropyran-4-yl)-(4-phenoxybenzenesulfonyl)amino]-propionic acid;  
 3-[(4-hydroxycarbamoyltetrahydropyran-4-yl)-(4-phenoxybenzenesulfonyl)amino]-propionic acid ethyl ester;  
 3-[(4-(4-fluorophenoxy)benzenesulfonyl)-(4-hydroxycarbamoylpiperidin-4-yl)-amino]propionic acid ethyl ester;  
 3-[(4-(4-chlorophenoxy)benzenesulfonyl)-(1-hydroxycarbamoyl-1-methylethyl)amino]-propionic acid;  
 3-[(4-(4-chlorophenoxy)benzenesulfonyl)-(1-hydroxycarbamoyl-1-methylethyl)amino]-propionic acid ethyl ester;  
 3-[(4-(4-fluorophenoxy)benzenesulfonyl)-(1-hydroxycarbamoylcyclohexyl)amino]-propionic acid;  
 3-[(1-hydroxycarbamoylcyclopentyl)-(4-phenoxybenzenesulfonyl) amino]propionic acid;  
 3-[(4-(4-chlorophenoxy)benzenesulfonyl)-(1-hydroxycarbamoylcyclopentyl)amino]-propionic acid

and pharmaceutically acceptable salts thereof.

13. The use of a prodrug compound of formula (Id):



wherein

X¹ is (C₁-C₆)alkyl;

X² is (C₁-C₆)alkyl; or

X¹ and X² taken together with the carbon atom to which they are attached form a ring selected from (C₅-C₇) cycloalkyl, 4-tetrahydropyranyl and 4-piperidinyl;

Y is a substituent on any of the carbon atoms of the phenyl ring capable of supporting an additional bond, preferably from 1 to 2 substituents (more preferably one substituent, most preferably one substituent in the 4-position) on the phenyl ring, independently selected from hydrogen, fluoro, chloro, trifluoromethyl, (C₁-C₆) alkoxy, trifluoromethoxy, difluoromethoxy and (C₁-C₆)alkyl; and

U and V are independently carbonyl, methylene, SO₂ or SO₃, and b is an integer from one to three wherein each methylene group is optionally substituted with hydroxy; in the manufacture of a medicament for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to Alzheimer's disease, stroke/cerebral ischemia, head trauma, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, migraine, cerebral amyloid angiopathy, AIDS, age-related cognitive decline, mild cognitive impairment and prion diseases.

14. A pharmaceutical composition useful in the treatment of a mammal afflicted with a disease, condition or disorder of the peripheral or central nervous system, including but not limited to Alzheimer's disease, stroke/cerebral ischemia, head trauma, spinal cord injury, multiple sclerosis, amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, migraine, cerebral amyloid angiopathy, AIDS, age-related cognitive decline, mild cognitive impairment and prion diseases, which comprises a compound of formula (I) according to Claim 1 and a suitable carrier or excipient.

15. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1, for administration to a mammal concomitantly receiving a non-steroidal anti-inflammatory drug.

16. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament combined with a non-steroidal anti-inflammatory drug for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1.

17. Use according to Claim 15 or 16 wherein the non-steroidal anti-inflammatory drug is selected from the group consisting of piroxicam, diclofenac, naproxen, flubiprofen, fenoprofen, ketoprofen and ibuprofen, fenamates, such as mefenamic acid, indomethacin, sulindac, apazone, and pyrazolones.
- 5 18. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1, for administration to a mammal concomitantly receiving a COX-1 or COX-2 inhibitor.
- 10 19. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament combined with a COX-1 or COX-2 inhibitor for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1.
- 15 20. Use according to Claim 18 and 19 wherein the COX-2 inhibitor is selected from the group consisting of celecoxib and rofecoxib.
- 20 21. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1, for administration to a mammal concomitantly receiving a CNS agent.
- 25 22. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament combined with a CNS agent for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1.
- 30 23. Use according to Claim 21 or 22 wherein the CNS agent is selected from the group consisting of an antidepressant selected from sertraline, fluoxetine, and paroxetine; an anti-Parkinsonian drug selected from deprenyl, L-dopa, and requip; miratex; MAOB inhibitors selected from selegine, rasagiline; COMP inhibitors and tolcapone; A-2 inhibitors; dopamine reuptake inhibitors; NMDA antagonists; nicotine agonists; dopamine agonists; inhibitors of neuronal nitric oxide synthase; anti-Alzheimer's drugs; acetylcholinesterase inhibitors selected from galantamine, metrifonate, donepezil, Exelon, and tetrahydroaminoacridine; COX-1 inhibitors, COX-2 inhibitors selected from celecoxib and rofecoxib; propentofylline; an anti-stroke medication; NR2B selective antagonists; glycine site antagonists; and a neutrophil inhibitory factor (NIF).
- 35 24. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1, for administration to a mammal concomitantly receiving an estrogen modulator.
- 40 25. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament combined with an estrogen modulator for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1.
- 45 26. Use according to claim 24 or 25 wherein the estrogen modulator is selected from the group consisting of estrogen, raloxifene, tamoxifene, droloxifene and lasofoxifene.
- 50 27. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1, for administration to a mammal concomitantly receiving an agent that results in a reduction of A $\beta$ 1-40/1-42.
- 55 28. The use of a compound of formula (I) according to Claim 1 in the manufacture of a medicament combined with an agent that results in a reduction of A $\beta$ 1-40/1-42 for the treatment in a mammal of a disease, condition or disorder of the peripheral or central nervous system, including but not limited to those described in Claim 1.
29. Use according to Claim 28 and 29 wherein the agent that results in reduction of A $\beta$ 1-40/1-42 is selected from the group consisting of amyloid aggregation inhibitors and secretase inhibitors.



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 00 30 8442

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl. 7)
X,P	WO 99 52889 A (PFIZER PRODUCTS INC.) 21 October 1999 (1999-10-21) * claims 1-11 * * page 7, line 32 - page 8, line 30 *	1,14-29	A61K31/18 A61P25/16 A61P25/28
X,D	WO 96 27583 A (PFIZER INC.) 12 September 1996 (1996-09-12) * claims 1-17 *	1-7,10, 11,14-29	
X,D	WO 98 33768 A (PFIZER PRODUCTS INC.) 6 August 1998 (1998-08-06) * claims 1-14 * * page 6, line 26 - page 7, line 2 *	1-6, 14-29	
X	WO 96 40101 A (CIBA-GEIGY AG) 19 December 1996 (1996-12-19) * page 20, line 13 - line 18 * * page 27, line 10 - line 21 * * claims 1-12 *	1-6, 14-19	
X	US 5 817 822 A (P. G. NANTERMET ET AL) 6 October 1998 (1998-10-06) * column 1, line 17 - column 2, line 11 * * column 17, line 54 - column 18, line 3 *	1-6, 14-29	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A61K
X	EP 0 895 988 A (PFIZER PRODUCTS INC.) 10 February 1999 (1999-02-10) * the whole document *	1-6, 14-29	
X,D	WO 99 07675 A (PFIZER PRODUCTS INC.) 18 February 1999 (1999-02-18) * claims 1-15 * * page 5, line 11 - line 16 *	14-29	
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 13 February 2001	Examiner Siatou, E
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention F : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1533 03 82 (PC/C01)



**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 8442

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

13-02-2001

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9952889	A	21-10-1999	AU 3271099 A	01-11-1999
			EP 1070058 A	24-01-2001
			HR 20000648 A	31-12-2000
			NO 20005076 A	07-12-2000
WO 9627583	A	12-09-1996	AU 707510 B	15-07-1999
			AU 5029396 A	23-09-1996
			BR 9607362 A	30-12-1997
			CA 2214720 A	12-09-1996
			CN 1181066 A	06-05-1998
			CZ 9702782 A	11-11-1998
			EP 0813520 A	29-12-1997
			FI 973613 A	05-11-1997
			HU 9800462 A	28-07-1998
			JP 11501910 T	16-02-1999
			NO 974103 A	05-11-1997
			NZ 303860 A	26-08-1998
			PL 322131 A	05-01-1998
			US 5863949 A	26-01-1999
WO 9833768	A	06-08-1998	AP 826 A	28-04-2000
			AU 721748 B	13-07-2000
			AU 8191798 A	25-08-1998
			BG 103616 A	29-02-2000
			BR 9807815 A	08-03-2000
			CN 1249743 T	05-04-2000
			EP 0977733 A	09-02-2000
			HR 980058 A	31-12-1998
			JP 2000508350 T	04-07-2000
			NO 993730 A	01-10-1999
			PL 335027 A	27-03-2000
WO 9640101	A	19-12-1996	US 5646167 A	08-07-1997
			AU 6124996 A	30-12-1996
			US 5817822 A	06-10-1998
US 5817822	A	06-10-1998	US 5646167 A	08-07-1997
			US 5552419 A	03-09-1996
			US 5506242 A	09-04-1996
			AU 6124996 A	30-12-1996
			WO 9640101 A	19-12-1996
			US 5672615 A	30-09-1997
			AT 196762 T	15-10-2000
			AU 692553 S	11-06-1998
			AU 2536995 A	19-01-1996
			CA 2192092 A	04-01-1996

EPO FORM P0455

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 8442

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

13-02-2001

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5817822	A		DE 69519024 D	09-11-2000
			EP 0766672 A	09-04-1997
			ES 2151599 T	01-01-2001
			FI 965156 A	20-12-1996
			HU 76548 A	29-09-1997
			WO 9600214 A	04-01-1996
			JP 11505502 T	21-05-1999
			NO 965568 A	17-02-1997
			ZA 9505206 A	27-12-1995
-----				
EP 895988	A	10-02-1999	BR 9803128 A	02-05-2000
			JP 11116549 A	27-04-1999
-----				
WO 9907675	A	18-02-1999	AU 8125398 A	01-03-1999
			BG 104118 A	29-09-2000
			BR 9811868 A	15-08-2000
			CN 1265646 T	06-09-2000
			EP 1003720 A	31-05-2000
			HR 980438 A	31-12-1999
			NO 20000604 A	07-02-2000
			PL 338633 A	06-11-2000
-----				

EPO FORM P0429

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82